arXiv:gr-qc/9912090v1 21 Dec 1999

K-causality and degenerate spacetimes

H.F.Dowker[a,1], R.S.Garcia[b,2], S.Surya[c,3]

[1] Dept. of Physics, Queen Mary and Westfield College, London E1 4NS, UK.
[2] D.A.M.T.P., University of Cambridge, Cambridge CB3 9EW, UK.
[3] TIFR, Homi Bhabha Rd, Mumbai 400 005, India.

February 7, 2008

Abstract

The causal relation K^+ was introduced by Sorkin and Woolgar to extend the standard causal analysis of C^2 spacetimes to those that are only C^0. Most of their results also hold true in the case of spacetimes with degeneracies. In this paper we seek to examine K^+ explicitly in the case of Lorentzian topology changing Morse spacetimes containing isolated degeneracies. We first demonstrate some interesting features of this relation in globally Lorentzian spacetimes. In particular, we show that K^+ is robust and that it coincides with the Seifert relation when the spacetime is stably causal. Moreover, the Hawking and Sachs characterisation of causal continuity translates into a natural expression in terms of K^+ for general spacetimes. We then examine K^+ in topology changing Morse spacetimes both with and without the degeneracies and find further characterisations of causal continuity.

[a]f.dowker@qmw.ac.uk, [b]R.Garcia@damtp.cam.ac.uk, [c]ssurya@tifr.res.in

1 Introduction

The causal relation K^+ was introduced in [1] to extend elements of standard global analysis concerning C^2 Lorentzian metrics to those which are only C^0. The main obstruction to such an extension within the framework of standard causal analysis is that the existence of the convex normal neighbourhoods is not guaranteed so that compactness of the space of causal curves cannot be proved in the usual way [2, 3]. The alternate causal relation K^+ was used in [1] to construct an appropriate compactness theorem in the C^0 case.

One of the original motivations of the work in [1] was to develop a causal analysis that would work in spacetimes with degeneracies. Indeed, while the usual chronological and causal relations are ill-defined at the degeneracies, K^+ can be naturally extended to cover these points as claimed in [1]. This present paper has grown out of our interest in the causal structure of causal Lorentzian topology changing spacetimes, which necessarily possess degeneracies [4, 5, 6, 7]. Traditionally, such degeneracies are excised from the spacetime, so that what remains is a globally Lorentzian spacetime. This was the procedure followed in [7, 8], to examine the causal continuity of Morse spacetimes. One way to incorporate the degeneracies in the overall causal structure would be to treat them as the boundaries of the spacetime, in the sense of [9]. This procedure, however, is not without ambiguities. Since K^+ can be defined in the degenerate spacetimes directly, it seems to be a promising alternative. In this paper, our main aim is to make sense of K^+ in topology changing Morse spacetimes which possess isolated degeneracies. Along the way, we demonstrate a number of interesting properties of K^+ in both globally Lorentzian spacetimes and those with isolated degeneracies.

An outline of our results is as follows. In the first section we briefly review some standard definitions of causal analysis before introducing the relation K^+. We then demonstrate that in a globally Lorentzian spacetime, K^+ is stable to certain changes in the chronological relation I^+ on which it is defined. For a C^2 globally Lorentzian spacetime (M, g), we show that K^+ changes minimally when we remove a set A of isolated points from M, namely, $K^+_{M-A} = K^+_M \cap (M - A) \times (M - A)$. K^+ thus appears to be "robust" to such changes. Next, we demonstrate that in a stably causal globally Lorentzian spacetime K^+ is precisely the Seifert relation J^+_S which provides us with an intuitive picture of K^+. We then show that in globally Lorentzian spacetimes, the Hawking and Sachs criterion for causal continuity [10] is equivalent to the condition, $K^\pm(x) = \overline{I^\pm(x)}$ for all x, which in turn is equivalent to the statement that (I^+, K^+) constitute a *causal structure*, as defined by Penrose and Kronheimer [11].

In the second section, we examine the special case of Morse spacetimes. When the degeneracies are excised, we show that in a Morse spacetimes for which the Morse function has only one critical value, $K^\pm(x)$ coincides with $\uparrow I^-(x)$ and $\downarrow I^-(x)$, respectively. The condition for causal continuity is shown to be equivalent to inner continuity of $\mathrm{Int}\,(K)^\pm(cdot)$. We then reinstate the degeneracies and show that again, K^+ is unaltered by changes in the chronological relation. We conjecture that the K^+'s in the

spacetimes with and without the degeneracies are related in a particular way and propose a definition of "k-causal continuity" in the unpunctured spacetime. We then show that the above conjecture implies the k-causal continuity in the spacetime with the degeneracies is equivalent to the causal continuity of the spacetime with the degeneracies excised. We also show that, independently of whether the conjecture holds, index 1 and $n-1$ Morse spacetimes are k-causally discontinuous, as expected [7, 8].

In the final section we speculate on the relationship between our work and the idea that spacetime is fundamentally discrete [12]. For example, the presence or absence of an isolated point in our continuum approximations should be irrelevant as far as any underlying discrete causal structure goes. Because of its apparent robustness, the relation K^+ seems a reasonable candidate to bridge the gap between continuous and discrete causal structures.

2 K^+ in globally Lorentzian spacetimes

We review some standard definitions of causal structure of spacetimes. In this section, (M, g) denotes a time orientable globally Lorentzian spacetime, with g of differentiability class C^0 unless otherwise specified.

A timelike or null vector v is future pointing if $g(v, u) < 0$ and past pointing if $g(v, u) > 0$, where u is a vector field that define the time orientation. A *future directed timelike path* in M is a C^1 function $\gamma : [0, 1] \to M$ whose tangent vector is future pointing timelike at $\gamma(t)$ for each $t \in [0, 1]$. Past directed paths are similarly defined using past pointing tangent vectors. We write $x <<_U y$ whenever there is a future directed timelike path γ with $\gamma(0) = x$ and $\gamma(1) = y$ totally contained in U. The image of a future (past) directed timelike path is a *future (past) directed timelike curve*. Given a point $x \in M$ we use the notation $I^+(x, U) \equiv \{y : x <<_U y\}$ for the chronological future of x relative to U. When $U = M$ we will often simplify to $x << y$ and $I^+(x) = I^+(x, M)$. The symbol $>>_U$ and the set $I^-(x, U)$ are defined dually. The chronos relation $I^+ \subset M \times M$ is defined by $I^+ \equiv \{(x, y) : x << y\}$. The chronology is an open subset of $M \times M$. As a relation, it is always transitive (i.e., $x << y$, $y << z \Rightarrow x << z$) and if the chronology condition holds, namely, if the spacetime contains no closed timelike curve, it is also antireflexive (i.e. there is no pair (x, y) with $x << y$ and $y << x$). The image of a future (past) directed causal path is a *future (past) directed causal curve*.

The *causal future* $J^+(x)$ and past $J^-(x)$ are similarly defined through causal paths with tangent vector timelike or null. One has $I^+(x) \subset J^+(x) \subset \overline{I^+}(x)$ and dual. The set of all pairs (x, y) with $x \in J^-(y)$ also defines a transitive relation in $M \times M$.

Given two metrics g_1, g_2 in a fixed manifold M, we write $g_2 > g_1$ if at every point $x \in M$, the g_2-lightcone is strictly larger than the g_1-lightcone. The *Seifert future* $J^+_S(x)$ of a point x in the spacetime (M, g) is defined as the intersection:

$$J^+_S(x) = \cap_{\bar{g}>g} J^+(x; \bar{g}) \tag{1}$$

It's not hard to see that $J^+(x) \subset J_S^+(x) = \overline{J_S^+(x)}$. Similarly one defines the Seifert past. It follows immediately from these definitions that $x \in J_S^-(y)$ iff $y \in J_S^+(x)$, therefore J_S^+ and J_S^- define the same relation in $M \times M$, which we simply call J_S.

We will need the *common future* $\uparrow U$ of an open set U, which is the interior of the set of all points connected to each point in U along a future directed timelike curve, i.e.,

$$\uparrow U \equiv \mathrm{Int}\left(\{x : \; x >> u \quad \forall u \in U\}\right) \tag{2}$$

It is clear from the definition that if $U \subset V$ then $\uparrow V \subset \uparrow U$. In addition the chronological future of a point is contained in the common future of its chronological past, namely $I^+(x) \subset \uparrow I^-(x)$; and a similar set inclusion holds for their duals, $I^+(x) \subset \uparrow I^-(x)$ (Proposition 1.1[10]).

Let F be a function which assigns to each event x in M an open set $F(x) \subset M$. Then we say that F is *inner continuous* if for any x and any compact set $C \subset F(x)$, there exists a neighbourhood U of x with $C \subset F(y)$ $\forall y \in U$. And F is said to be *outer continuous* if for any x and any compact set $C \subset M - \overline{F(x)}$, there exists a neighbourhood U of x with $C \subset M - \overline{F(y)}$ $\forall y \in U$. The chronological past and future are inner continuous in any globally Lorentzian spacetime [10].

Definition 1 *A time-orientable distinguishing spacetime, (M, g) is said to be causally continuous if it satisfies any of the following equivalent conditions [10]:*

(I) for all events x in M we have $I^+(x) = \uparrow I^-(x)$ and $I^-(x) = \downarrow I^+(x)$.

(II) for all events x and y, $x \in \overline{I^+}(y)$ iff $y \in \overline{I^-}(x)$;

(III) for all events $x \in M$, $I^+(x)$ and $I^-(x)$ are outer continuous.

(IV) for all events $x \in M$, $J_S^+(x) = \overline{I^+(x)}$ and $J_S^-(x) = \overline{I^-(x)}$

Causality holds in a subset S of M if there are no causal loops based at points in S. *Strong causality* holds in a subset S of M if for every point $s \in S$ any neighbourhood U of s contains another neighbourhood V of s that no causal curve intersects more than once. A spacetime (M, g) is said to be *stably causal* if there is a metric g', whose lightcones are strictly broader than those of g and for which the spacetime (M, g') is causal. A spacetime is stably causal if and only if it admits a global time function, i.e., a function which is monotonically increasing along every future directed causal curve. The spacetime M is said to be *causally simple* if $J^{\pm}(x) = \overline{I^{\pm}(x)}$ for every point $x \in M$. A spacetime is *globally hyperbolic* if it possesses a spacelike hypersurface which every inextendiblecausal curve intersects exactly once.

In [11] a *causal structure* is defined axiomatically as follows.

Definition 2 *Given a set X and three relations, $<<$, $<$ and $\rightarrow$, on X, the quadruple $(X, <<, <, \rightarrow)$ is said to define a causal structure if the following conditions are satisfied for every x, y and z in X:*

$$
\begin{array}{rl}
(i) & x < x \\
(ii) & \textit{if} \quad x < y \quad \textit{and} \quad y < z, \quad \textit{then} \quad x < z \\
(iii) & \textit{if} \quad x < y \quad \textit{and} \quad y < x, \quad \textit{then} \quad x = y \\
(iv) & \textit{not} \quad x << x \\
(v) & \textit{if} \quad x << y, \quad \textit{then} \quad x < y \\
(vi+) & \textit{if} \quad x < y \quad \textit{and} \quad y << z, \quad \textit{then} \quad x << z \\
(vi-) & \textit{if} \quad x << y \quad \textit{and} \quad y < z, \quad \textit{then} \quad x << z \\
(vii) & x \rightarrow y \quad \textit{iff} \quad x < y \quad \textit{and not} \quad x << y
\end{array}
$$

Note that in any globally Lorentzian spacetime, with $x < y$ denoting $x \in J^-(y)$, $x << y$ denoting $x \in I^-(y)$ and $x \rightarrow y$ denoting $x \in J^-(y) - I^-(y)$, all the axioms above are automatically satisfied, except for (iii), which is the condition of causality.

In [1] the relation K^+ is defined from I^+ as follows:

Definition 3 *Given a spacetime (M, g) the relation K^+ on $M \times M$ is the smallest transitive relation that contains I^+ and is closed as a subset of $M \times M$.*

K^+ is thus the *transitive closure* of I^+ in $M \times M$. By analogy with I^+, we can define $K^+(x) = \{y : \ (x, y) \in K^+\}$ and write $x \prec y$ if $y \in K^+(x)$. $K^-(x)$ is defined similarly. We also define $K(x, y) = K^+(x) \cap K^-(y)$. Further, if $U \subset M$, then the relation K^+_U is defined with respect to I^+_U, the chronology relation in $U \times U$. We will use either $y \in K^+(x, U)$ or $x \prec_U y$ to denote a pair (x, y) in K^+_U.



Figure 1: This is a region of the spacetime obtained by removing the semiline $t = 0$, $x \leq 0$ from 2-dimensional Minkowski space. For the point q we have $J^+(q) = \overline{I^+(q)} \subset K^+(q)$, where the inclusion is strict since $s \in K^+(q) - \overline{I^+(q)}$; while for r we have $J^+(r) \subset \overline{I^+(r)} = K^+(r)$, and the inclusion here is also strict, since $s \in \overline{I^+}(r) - J^+(r)$

We note that $\overline{I^+} \subseteq K^+$, where $\overline{I^+}$ denotes the closure of I^+ in $M \times M$ and $K^+(x)$ and $K^-(x)$ are closed $\forall x \in M$. Also, $J^+(x) \subseteq \overline{I^+(x)} \subseteq K^+(x)$. A simple example, from

[1], where both set-inclusions are strict is 2-dimensional Minkowski spacetime with the $t = 0$, $x < 0$ semiline removed (see figure 1). For any compact set $C \subset M$ the set $K^+(C)$ is closed and $\overline{\mathrm{Int}\,((K^{\pm}(x)))} = K^{\pm}(x)$. Thus we obtain the following,

Lemma 1 *The functions* $\mathrm{Int}\,(K^+(\cdot))$ *and* $\mathrm{Int}\,(K^-(\cdot))$ *are outer continuous.*

Proof Given a point x and a compact set $C \subset M$ with $C \subset M - \overline{\mathrm{Int}\,(K^+(x))} = M - K^+(x)$, $x \notin K^-(C)$ and therefore, by closure of $K^-(C)$, there must be a neighbourhood U_x with $U_x \cap K^-(C) = \emptyset$ $\Box$.

Now, $U \subset M$ is said to be *K-causal* if $x \prec y$ and $y \prec x$ simultaneously imply $x = y$ for all points $x, y \in U$. An open set $O \subset M$ is said to be *K-convex* if for every pair $x, y \in O$ with $x \prec y$, $K(x, y) \subset O$. A K-convex set O is further *K-globally hyperbolic* if for every pair of points $x, y \in O$ the interval $K(x, y)$ is compact.

In contrast with I^+ and J^+, the relation K^+ is defined without direct reference to curves in the spacetime. As a consequence, certain properties that are obvious for the relation J^+, need to be established for K^+. The properties we list here have been proved in [1], and we mark these with an "(SW)":

Lemma 2 (SW) *Given two open sets $U \subset O$ in M and two points $x, y \in M$ then: (i) if $x \prec_U y$, then also $x \prec_O y$; (ii) if $x \prec_O y$ and U is K-convex with respect to K^+_O, then also $x \prec_U y$.*

Lemma 3 (SW) *Let $B \subset M$ be set with compact boundary ∂B and suppose there are two points $x \in B$ and $y \notin B$ with $x \prec y$, then there must be a point $z \in \partial B$ such that $x \prec z \prec y$.*

Lemma 4 (SW) *Let $B \subset M$ be an open set with compact closure and the points $x, y \in B$ have $x \prec y$ but not $x \prec_B y$, then there is a point $z \in \partial B$ such that $x \prec z \prec y$.*

Lemma 5 (SW) *If (M, g) is K-causal, then every point $x \in M$ has arbitrarily small K-convex neighbourhoods.*

Lemma 6 (SW) *In a C^2 Lorentzian spacetime $J^+(x) = K^+(x)$ for every x if and only if the spacetime is causally simple.*

The significance of lemma 5 (lemma 16 in [1]) in the context of globally Lorentzian spacetimes is that strong K-causality and K-causality are equivalent [1]. This implies, in particular, that if a spacetime is J-causal, but not strongly J-causal, then it is not K-causal. Indeed, take a causal spacetime with a strong causality violation as described in [3], namely there exists a pair of points $x \neq y$ satisfying (i) $y \in J^+(x)$ and (ii) $I^-(y) \subset \downarrow I^+(x)$. (i) $\Rightarrow x \prec y$ and (ii) along with the closure of $K^+ \Rightarrow y \prec x$, i.e., the spacetime is not K-causal. Lemma 5 leads to another intuitive result,

Corollary 1 (SW) *If the C^2 spacetime (M, g) is K-causal, then every point $x \in M$ has a neighbourhood U such that $K^+_{|U} = J^+_{|U}$.*

Indeed, let U be a K-convex neighbourhood contained in a J-globally hyperbolic neighbourhood V of x, for which $K^+_V = J^+_V$. Then $K^+_{|U} = K^+_U = J^+_U = J^+_{|U}$, where the last set-equality follows because K-convexity implies J-convexity.

Note that not all the properties of K^+ given above are valid in degenerate spacetimes. Apart from the obvious failure, namely, lemma 6, lemma 5 too is not readily available, since the Lorentzian character of g is used at every point in its proof [1]. However, lemmas 2, 3 and 4 are valid, since their proofs do not depend on the assumption that the metric g is everywhere Lorentzian or even that timelike curves exist through every point of the spacetime.

2.1 The relation K^+ and the Seifert relation

Intuitively, the Seifert relation, J^+_S, depends on the stability of the lightcones against perturbations. We show that, indeed, in globally Lorentzian causally stable spacetimes, K^+ and J^+_S coincide.

Proposition 1 *In a globally Lorentzian spacetime $K^+ \subseteq J^+_S$. If the spacetime is stably causal then $K^+ = J^+_S$.*

Proof Clearly $I^+ \subset J^+_S$. Thus, in order to show that $K^+ \subseteq J^+_S$, we need to prove closure and transitivity of J^+_S. To prove the equivalence for stably causal spacetimes, it then suffices to show minimality of J^+_S. For this we are assisted by a lemma proved by Seifert under the assumption of stable causality [13, 10].

(i) (a) We first prove that J^+_S is closed. Consider a pair $(x, y) \notin J^+_S$. Since $J^-_S(y)$ is closed [10] in M, there exists a neighbourhood V_x of x disjoint from $J^-_S(y)$. Pick a point x' in $V_x \cap I^-(x)$. Again, since $J^+_S(x')$ is closed, there is a neighbourhood V_y disjoint from $J^+_S(x')$. Take a point $y' \in V_y \cap I^+(y)$. There is a metric $g' > g$ so that $y' \notin J^+(x', g')$. Consider the new neighbourhoods $U_x \subset I^+(x', g')$ and $U_y \subset I^-(y', g')$ respectively, of, x and y. For any pair of points $(z, w) \in U_x \times U_y$ we must have $z \notin J^-(w, g')$, since

otherwise we would deduce that $y' \in I^+(x', g')$. Therefore $U_x \times U_y$ contains no pair in J_S^+.

(b) Next we prove that J_S^+ is transitive. Consider $y \in J_S^+(x)$ and $z \in J_S^+(y)$, so that $z \in J^+(y, g')$ and $y \in J^+(x, g')$ $\forall g' > g$. By transitivity of $J^+(\cdot, g')$ this implies that $z \in J^+(x, g')$.

(ii) Lemma 1 in [13] says that in a stably causal spacetime J_S^+ is a partial order. Lemma 2 in [13] then states that if J_S^+ is a partial order then it is contained in any relation $\mathcal{J}^+$ satisfying $J^+(x) \subset \overline{\mathcal{J}^+(x)} = \mathcal{J}^+(x)$. Since this is satisfied by the relation K^+, it follows that $J_S^+ \subset K^+$. $\Box$

2.2 Robustness of K^+

As stated in the introduction, we would like to introduce K^+ as the natural causal relation in a spacetime with degeneracies, specifically topology changing spacetimes. In previous work we have analysed such topology changing spacetimes by excising these degeneracies and using the standard definitions of chronology (I^+) and causality (J^+) as defined above. One might wonder whether the causal properties of a spacetime depend crucially on the absence of the excised point and whether adding it back in in some way mightn't make a big difference.

As a first step towards justifying the use of K^+ we will show that, at least in the case of a point excised from and then added back into a globally Lorentzian spacetime, the causal structure given by K^+ is as robust as can be imagined and is only minimally altered by the excising and adding process, no matter how it is performed. This seems physically reasonable too as we mentioned before.

We first prove a lemma which is clearly related to the issue of missing points and which we will use later.

Lemma 7 *Let (M, g) be a globally Lorentzian, C^2 spacetime and $p \in M$. Then $y \in I^+(x)$, $x, y \neq p$, implies that $y \in I^+(x, M - p)$.*

Proof There exists a future directed timelike path $\gamma : [0, 1] \to M$ so that $\gamma(0) = x$ and $\gamma(1) = y$. Let Γ be its image in M. Either Γ does not intersect p, in which case we are done, or it does. Suppose the latter. Let t_1 be the first value of t for which $\gamma(t) = p$ and let $t_2 \equiv \sup\{t \in [0, 1] : \gamma(t) = p\}$.[1]

By continuity $t_1 \neq 0$, $t_2 \neq 1$ and $\gamma(t_2) = p$. Consider therefore the new path $\widetilde{\gamma} : [0, 1] \to M$ formed by joining the two pieces of γ on $[0, t_1]$ and $[t_2, 1]$. Its image $\widetilde{\Gamma}$ intersects p exactly once.

[1] We do not think it possible that γ intersect p infinitely often but it is easier here to use the supremum construction rather than prove this.

Now consider a convex normal neighbourhood of p, U. Choose a point $y' \in U \cap \widetilde{\Gamma} \cap I^+(p)$ such that all points on $\widetilde{\Gamma}$ between p and y' lie in U. Similarly choose a point $x' \in U \cap \widetilde{\Gamma} \cap I^-(p)$ such that all the points on $\widetilde{\Gamma}$ between x' and p lie in U. Define a new neighbourhood of p, $V = U \cap I^-(y', U) \cap I^+(x', U)$. Choose $p' \in V$ such that there are no timelike curves in U between p' and p. Such a p' is guaranteed to exist because U is a convex normal neighbourhood which, roughly speaking, has the causal structure of a convex open set in Minkowski spacetime.

By construction there is a timelike path γ_x from x' to p' and one, γ_y from p' to y'. Now we build a new timelike path γ' from x to y by joining $\widetilde{\gamma}$ from x to x', γ_x from x' to p', γ_y from p' to y' and $\widetilde{\gamma}$ from y' to y. By construction, its image Γ' cannot intersect p. Hence the result. □

A simple corollary is that a timelike curve can be deformed to avoid any finite number of points whilst remaining timelike.

Consider the C^2 spacetime (M, g). Let $A \subset M$ be a set of finitely many points. On M we can consider a number of different relations,

$$I^+_M \equiv I^+ \tag{3}$$
$$\widetilde{I}^+_M \equiv I^+_M \cap (M - A) \times (M - A) \tag{4}$$
$$\widehat{I}^+_M \equiv \{(x, y) : x, y \notin A, x \in I^-(y, M - A)\} \tag{5}$$

By lemma 7 we have $\widetilde{I}^+ = \widehat{I}^+$. So we can consider the following, potentially different, relations: K^+_M the transitive closure of I^+_M and $\widetilde{K}^+_M$ the transitive closure of $\widetilde{I}^+_M$. The following lemma shows that these are equal.

Lemma 8 *Let (M, g), $A \subset M$, K^+_M and $\widetilde{K}^+_M$ be as above. Then $K^+_M = \widetilde{K}^+_M$.*

Proof Since it is minimal, K^+_M is the transitive closure of any relation whose closure is equal to the closure of I^+_M so it suffices to prove that $\overline{I^+_M} = \overline{\widetilde{I}^+}$. We have $\overline{\widetilde{I}^+} \subset \overline{I}^+$. Consider $(x, y) \in \overline{I^+}$, so that there exists a sequence of points in I^+, $(x_k, y_k) \to (x, y)$. There are four cases depending on whether x and/or y is in A. We will prove two as the remaining cases are similar. (i) If $x, y \notin A$ then $\exists N > 0$ such that $k > N$ implies that $x_k, y_k \notin A$ and so $(x, y) \in \overline{\widetilde{I}^+}$. (ii) If $x \in A$ and $y \notin A$ then either $\exists\ N > 0$ such that $x_k = x\ \forall k > N$, or there's an infinite subsequence $x_{k'} \to x$ and $x_{k'} \notin A$. In the latter case we're done as in (i). In the former case we have $(x, y_k) \in I^+$. Choose a sequence of points $x_k \in I^-(x)$i, $x_k \neq x$, which converge to x and we are done as before. □

On the punctured spacetime $N \equiv M - A$ we have the relations $I^+_N \equiv \widehat{I}^+_M$ and $\widetilde{I}^+_N \equiv \widetilde{I}^+_M$. These two relations are equal as before. The possible corresponding causal relations on N are K^+_N the transitive closure of I^+_N and $\breve{K}^+_N \equiv K^+_M \cap N \cap N$.

The next result shows that if M is K-causal $K^+_N = \breve{K}^+_N$, in other words, the absence of a finite $A \subset M$ alters the K^+ relation only to the extent of removing relations involving

points in A. Our proof is based on the set theoretic technique of transfinite induction, which is a generalisation of ordinary induction, to deal with constructions that do not occur through a countable number of steps.

Transfinite induction is used to verify that a certain property, call it X, satisfied by pairs in I^+ is preserved as new pairs are successively added in the construction of K^+, because of the requirements of closure and transitivity [1]. More concretely, one considers at an intermediate stage, the relation, $K^+_\alpha \subset K^+$ and the hypothesis, H_α, that X is satisfied by all points in K^+_α. To prove this hypothesis, one must first establish that H_0 is true, namely that X is satisfied by $K^+_0 \equiv I^+$. Next, given that X is satisfied for all K^+_β, where $\beta < \alpha$, one needs to show that it is true for α. At this stage, there are two cases that arise in this transfinite version of induction. If α is a limit ordinal the conclusion follows since $x \prec^\alpha y \Rightarrow x \prec^\beta y$ for $\beta < \alpha$. If α is not a limit ordinal, then $\exists \beta$ such that $\alpha = \beta + 1$, namely, $K^+_\alpha = K^+_\beta \cup (x, y)$. The new pair (x, y) has been added either because there are sequences $x_k \to x$ and $y_k \to y$ with $(x_k, y_k) \in K^+_\beta \; \forall \, k$ or because there is a point z such that $(x, z) \in K^+_\beta$ and $(z, y) \in K^+_\beta$. The essence of the proof is to show that in either case (x, y) also satisfies the property X. For more details, and to appreciate the difference between ordinary induction and transfinite induction, see [1, 14].

Proposition 2 *Let (M, g), $A \subset M$, $N \equiv M - A$, K^+_N and $\breve{K}^+_N$ be as above and let (M, g) be in addition K-causal. Then $\breve{K}^+_N = K^+_N$.*

Proof We will adopt the convention in this proof that any relation, I^+, K^+, $\prec$, *etc.*, that has no subscript labeling which manifold it is on will be understood to be on M.

First assume that A contains a single point p, so that $N = M - p$. $\breve{K}^+_N$ defines a closed transitive relation on N which includes I^+_N. Thus, all we need to show is that $\breve{K}^+_N \subset K^+_N$ since K^+_N is minimal. In other words, if a pair $(x, y) \in N \times N$ is such that $x \prec y$, we want $x \prec_N y$. It is convenient to establish this property by dividing the pairs (x, y) into two sets, namely (i) $p \notin K(x, y)$ and (ii) $p \in K(x, y)$.

(i) Here, we employ the methods of transfinite induction discussed above. Let H_α be the hypothesis that at stage α, if the pairs (x, y) satisfy $x \prec^\alpha y$ and $x, y \in N$, with $p \notin K(x, y)$, then $x \prec_N y$.

This is true at the initial step $K^+_0 \equiv I^+$, since $p \notin K(x, y) \Rightarrow p \notin I(x, y)$, so that any timelike curve from x to y must be contained in N. Now, consider the stage α and let H_β hold for every $\beta < \alpha$. If α is a limit ordinal, H_α follows. Otherwise, if $\alpha = \beta + 1$, we can assume that the pair added in going from β to α, is (x, y), and $p \in K(x, y) \Rightarrow H_\alpha$ is trivially satisfied, so that we only need to consider the case, $p \notin K(x, y)$.

First, suppose the new pair has been added because of closure, i.e., there are sequences $x_k \to x$ and $y_k \to y$ with $x_k \prec^\beta y_k$. The x_k, y_k are in N for large enough k. Moreover, by closure of K^+, we must since $p \notin K(x, y)$, for sufficiently large k, $p \notin K(x_k, y_k)$. In

other words, there is an integer m such that the hypothesis H_β applies to the tail of the sequences $x'_k = x_{k+m}$ and $y'_k = y_{k+m}$, so that $x'_k \prec_N y'_k$. Thus, by closure of K_N^+, $x \prec_N y$.

If the new pair has been added because of transitivity, then there is a point z with $x \prec^\beta z \prec^\beta y$. Since $p \notin K(x,y)$, $z \neq p$, and for the same reason $p \notin K(x,z)$ and $p \notin K(z,y)$. It follows from H_β that that $x \prec_N z \prec_N y$ and, by transitivity of K_N^+, we see that $x \prec_N y$.

Thus, we have shown that for all $(x,y) \in N$ satisfying $(x,y) \in K^+$ and $p \notin K(x,y)$, $x \prec_N y$. Note that we do not require the C^2 condition for this part of the proof.

(ii) Now we prove the property for the pairs (x,y), such that $p \in K(x,y)$. Let U be a K-convex neighbourhood such that $K_U^+ = \overline{I_U^+}$, which exists by virtue of corollary 1. [2] Choose $p_1, p_2 \in U$ such that $x \prec p_1 \prec p$ and $p \prec p_2 \prec y$. That such p_1 and p_2 exist is guaranteed by applying lemma 3 to the compact boundary of a subneighbourhood $V \subset \overline{V} \subset U$. Now since $K_{|U}^+ = K_U^+ = \overline{I_{|U}^+}$, we have $p \in \overline{I^+(p_1)}$ and $p_2 \in \overline{I^+(p)}$. We claim (see below) that this implies $p_2 \in \overline{I^+(p_1)}$, which means, by lemma 7, $p_2 \in \overline{I_N^+(p_1)} \subset K_N^+(p_1)$. Also, by K-causality we have $p \notin K(x,p_1)$ so that part (i) implies $x \prec_N p_1$. Similarly $p_2 \prec_N y$. Thus we have $x \prec_N p_1 \prec_N p_2 \prec_N y$, and since K_N^+ is transitive we conclude $x \prec_N y$.

If A contains more than one point, note that because $K^+(x, M-p_1) = K^+(x) \cap (M-p_1)$, for any $p_1 \in M$, it follows that

$$\begin{aligned} K^+\left(x, M-(p_1 \cup p_2)\right) &= K^+(x, M-p_1) \cap ((M-p_1)-p_2) \\ &= \left(K^+(x) \cap (M-p_1)\right) \cap (M-(p_1 \cup p_2)) \\ &= K^+(x) \cap (M-(p_1 \cup p_2)), \end{aligned} \tag{6}$$

for any $p_2 \in M$ with $p_2 \neq p_1$. □

It remains to prove the claim:

Claim 1 *If $z \in \overline{I^+(y)}$ and $y \in \overline{I^+(x)}$ then $z \in \overline{I^+(x)}$.*

Proof There exist sequences $z_k \to z$ and $y_l \to y$ with $z_k \in I^+(y)$ and $y_l \in I^+(x)$. Consider $I^-(z_k)$ for each k. There's a neighbourhood of y contained in $I^-(z_k)$ and so there's an l_k such that $y_{l_k} \in I^-(z_k)$. Hence $z_k \in I^+(x)$. □

For any finite set A in a K-causal spacetime, (M,g), the above result makes it easy to show that the causal future and past of the points in A are determined by the relation K_N^+ on $N \equiv M - A$ simply by taking the closure in $M \times M$, *i.e.*, $K_M^+ = \overline{K_N^+}$.

Lemma 9 *Let (M,g), A, N, K_N^+ and K_M^+ be as in proposition 2. Then $K_M^+ = \overline{K_N^+}$, where the closure is taken in $M \times M$.*

[2] Note that the C^2 requirement makes its appearance at this crucial stage in the proof.

Proof We know from proposition 2 that $K_N^+ = K_M^+ \cap N \times N$. So we have $\overline{K_N^+} \subset K_M^+$. On the other hand suppose $(x, y) \in K_M^+$. Then (i) $(x, y) \in A \times A$, or (ii) $(x, y) \in A \times N$, or (iii) $(x, y) \in N \times A$, or (iv) $(x, y) \in N \times N$. In each case there exists a sequence in K_N^+ which converges to (x, y). We will show this explicitly in case (i) only, as the other cases are similar and simpler. Choose a sequence of points $x_k \to x$, $x_k \in I^-(x) \cap N$. Similarly choose a sequence $y_k \to y$, $y_k \in I^+(y) \cap N$. Then (x_k, y_k) is a sequence that converges to (x, y) and $(x_k, y_k) \in K_N^+$. So $(x, y) \in \overline{K_N^+}$ and hence result. □

Another natural way to construct the causal relation on M from knowledge of N would be to consider I_N^+, the usual chronological relation on N, as a relation on M, and construct the transitive closure of it in $M \times M$. In other words take the transitive closure of what we called $\widehat{I}_M^+$. That this gives K_M^+ is the statement of lemma 8.

We conclude that the relation K^+ is robust against the subtracting and adding of isolated points.

2.3 Causal continuity in terms of the relation K

The definition of causal continuity due to [10] concerns globally Lorentzian spacetimes. However, causal continuity appears to be an important property in the study of topology changing Morse spacetimes, which contain isolated degeneracies [7]. In order to extend the notion of causal continuity to spacetimes with degeneracies, we seek alternative characterisations that involve K^+ alone, since, as we shall see shortly, this can be defined at the degeneracies.

First we prove that two conditions based on the pair (I^+, K^+) are equivalent to the standard definition.

Proposition 3 *In a K-causal spacetime (M, g) the following three conditions are equivalent.*

(A) The spacetime (M,g) is causally continuous.

(B) For every $x \in M$, $K^\pm(x) = \overline{I^\pm(x)}$.

(C) The relations (I^+, K^+) define a causal structure in M.

Proof We use the definition 1.II of causal continuity, namely, for any $x, y \in M$, $x \in \overline{I^-(y)} \Leftrightarrow y \in \overline{I^+(x)}$.

(A) implies (B). When causal continuity holds the subsets $\{(x, y) : y \in \overline{I^+(x)}\}$ and $\{(x, y) : x \in \overline{I^-(y)}\}$ are identical in $M \times M$ and hence define the same relation. This relation is transitive by claim 1 and contains I^+ in a minimal way, so we only need prove that it is closed in $M \times M$.

Suppose there are sequences $x_k \to x$ and $y_k \to y$ with $y_k \in \overline{I^+(x_k)}$. We show that causal

continuity implies that $y \in \overline{I^+(x)}$. For any $x' << x$, there is an integer k_0 such that $x_k \in I^+(x') \quad \forall \quad k > k_0$. Therefore we have a sequence $\{y_k\} \quad k > k_0$ with $y_k \in \overline{I^+(x')}$, which is a closed set in M, hence $y \in \overline{I^+(x')}$ and, by causal continuity, $x' \in \overline{I^-(y)}$. Since there is such an x' in every neighbourhood of x we conclude, by closure of $\overline{I^-(y)}$ in M, that $x \in \overline{I^-(y)}$, or equivalently $y \in \overline{I^+(x)}$.

(B) implies (A). If $K^\pm(x) = \overline{I^\pm(x)}$, then for every pair of points x, y we have $x \in \overline{I^-(y)}$ iff $x \in K^-(y)$ iff $y \in K^+(x)$ iff $y \in \overline{I^+(x)}$.

(B) implies (C). All but axioms (ii) and (vi) in the definition of a causal structure 2 are satisfied by (I^+, K^+) in any globally Lorentzian spacetime. Axiom (ii) is guaranteed by our requirement of K-causality, so we only need to consider (vi). Consider $x \prec y << z$. There exists neighbourhood U_y of y, such that $U_y \subset I^-(z)$. Now, $y \in K^+(x) = \overline{I^+(x)}$, so that there is a point $y' \in U_y$ with $y' \in I^+(x)$. Hence, by transitivity of I^+, $x << z$. Similarly for (vi-).

(C) implies (B). Let $x \prec y << z \Rightarrow x << z$ for all $x, y, z \in M$. i We have $\overline{I^+(x)} \subset K^+(x)$. Let us assume that there is an $x \in M$ for which $K^+(x) - \overline{I^+(x)} \neq \emptyset$. If every y in $K^+(x) - \overline{I^+(x)}$ is in the boundary of $K^+(x)$, we would have $\mathrm{Int}\,(K^+(x)) \subset \overline{I^+(x)}$ and therefore, from the paragraph before lemma 1, $K^+(x) \subset \overline{I^+(x)}$, thus contradicting the assumption. This means that there is a point y with a neighbourhood U_y contained in $K^+(x) - \overline{I^+(x)}$. Hence for any $z >> y$ in U_y we have $x \prec y << z$ and not $x << z$, which is again, a contradiction. □

Notice that "(A) implies (B)" can be immediately inferred from definition 1(IV) and theorem 1. We have nevertheless given a proof in terms of K^+ because it is simple enough and avoids the mention of J^+_S.

We now propose two characterisations of causal continuity involving only K^+. In the next section, we show that these characterisations are valid for Morse spacetimes, while in general, we can partly prove their equivalence to the conditions in proposition 3. In a K-causal spacetime (M, g) consider the conditions:

(D) $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$ are inner continuous.

(E) For any $x \in M$ and K-convex neighbourhood N of x, $\mathrm{Int}\,(K^+_N(x))$ and $\mathrm{Int}\,(K^-_N(x))$ are connected.

We have put forward (D) in analogy to definition 1(III). Remember that in any spacetime $I^+(\cdot)$ and $I^-(\cdot)$ are inner continuous, while $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$ are outer continuous. Since outer continuity of $I^\pm(\cdot)$ is one characterisation of causal continuity, it seems reasonable to expect that so must be inner continuity of $\mathrm{Int}\,(K^\pm(\cdot))$. This "duality" stems from the fact that I^+ is open as a relation in $M \times M$, while K^+ is closed.

For example, let us look at the past $I^-(\cdot)$. For any point x in a spacetime, $I^-(x) - I^-(y)$ becomes arbitrarily small as y approaches x from its past —this is the inner continuity of $I^-(\cdot)$—, while $K^-(z) - K^-(x)$ becomes arbitrarily small as z approaches

x from its future —outer continuity of $\mathrm{Int}\,(K^-(\cdot))$. An abrupt change of the past occurs at a point x if $I^-(z) - I^-(x)$ contains a fixed open set as z approaches x from its future, or, as we propose, if $K^-(x) - K^-(y)$ contains a fixed open set as y approaches x from its past[3].

As for (E), we have arrived at this condition by looking at different causally discontinuous spacetimes, all of which have in common a local disconnectedness of the K-pasts or K-futures. The details in the condition are as follows. We require N to be convex in order to avoid any spurious disconnectedness associated with the boundary of N (figure 2(a)). We impose the condition to hold for arbitrary N because often the past or future are disconnected only locally and become connected again in large enough neighbourhoods (figure 2(b)). Finally we express it in terms of $\mathrm{Int}\,(K^\pm(\cdot))$ rather than $K^\pm(\cdot)$ itself so that our condition extends to degenerate spacetimes: degeneracies will sometimes connect otherwise disconnected pasts and futures, but not their interiors (figure 2(c)).



Figure 2: In (a) we show a non-convex neighbourhood N of 2-dimensional Minkowski spacetime. For the point $x \in N$, $\mathrm{Int}\,(K^+(x, N))$ is disconnected. (b) is a picture of the spacetime obtained from 2-dimensional Minkowski by removing a spacelike segment (in bold). We show how the K-past of q, which is disconnected in N_1 (bounded by dashed lines), becomes again connected in N_2.

Lemma 10 *If the K-causal spacetime (M, g) is causally continuous, then*

(i) $\mathrm{Int}\,(K^+(\cdot))$ *and* $\mathrm{Int}\,(K^-(\cdot))$ *are inner continuous. (ii) For any $x \in M$ and K-convex neighbourhood N of x,* $\mathrm{Int}\,(K^+_N(x))$ *and* $\mathrm{Int}\,(K^-_N(x))$ *are connected.*

Proof (i) is immediate, since from proposition 3, we know that causal continuity implies that $\mathrm{Int}\,(K^+(x)) = \mathrm{Int}\,(\overline{I^+(x)}) = I^+(x)$, which is inner continuous [10].

For (ii), suppose that causal continuity holds, so that $K^\pm(x) = \overline{I^\pm(x)}$ for each x, and let N be K-convex, we show for example that $K^+_N(x) = \overline{I^+_N(x)}$. This suffices, since $I^+_N(x)$ is

[3]Because inner continuity of the past at x means roughly that it converges from "below" and outer continuity, that it converges from "above", one may envisage unifying the two conditions in a single notion of continuity, using an appropriate topology. It has been suggested [15] that the Vietoris topology [1] may be suitable for this purpose.

clearly connected. Now any $y \in K_N^+(x)$ is also in $K^+(x)$ and therefore in $\overline{I^+(x)}$. Hence there exists a sequence of points $y_k \to y$ with $y_k \in I^+(x)$ and we can assume that all the y_k lie in N, since this is an open set. Let Γ_k be a sequence of timelike curves from x to y_k. Then $\Gamma_k \subset K(x, y_k)$, which by convexity of N implies $\Gamma_k \subset N$. Therefore $y \in \overline{I_N^+(x)}$, as desired. $\Box$

With this we have shown that causal continuity implies conditions (D) and (E) above. Later we show that in Morse spacetimes (D) implies causal continuity. The following lemma provides a convenient way of expressing condition (D).

Lemma 11 *In a K-causal spacetime (M, g), $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$ are inner continuous if and only if for every $x, y \in M$, $x \in \mathrm{Int}\,(K^-(y)) \;\Leftrightarrow\; y \in \mathrm{Int}\,(K^+(x))$.*

Proof Suppose $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$ are inner continuous. If $x \in \mathrm{Int}\,(K^-(y))$, then there must be a neighbourhood U_y of y such that $x \in \mathrm{Int}\,(K^-(y'))\ \forall y' \in U_y$. Therefore $U_y \subset K^+(x)$ and $y \in \mathrm{Int}\,(K^+(x))$. Similarly $y \in \mathrm{Int}\,(K^+(x))$ implies $x \in \mathrm{Int}\,(K^-(y))$.

Conversely, suppose that for every $x, y \in M$, $x \in \mathrm{Int}\,(K^-(y)) \;\Leftrightarrow\; y \in \mathrm{Int}\,(K^+(x))$. Consider any $y \in M$ and the compact set $C \subset \mathrm{Int}\,(K^-(y))$. For every $x \in C$, the condition implies that we can find points $z >> x$ and $w << y$ such that $z \in \mathrm{Int}\,(K^-(w))$ and therefore neighbourhoods $U_x \subset I^-(z)$ of x and $U_y^x \subset I^+(w)$ of y so that $U_x \subset \mathrm{Int}\,(K^-(y'))\quad \forall y' \in U_y^x$. The cover $\{U_x :\ x \in C\}$ of C must have a finite subcover, so $C \subset \cup_{j=1}^N U_{x_j}$. Let $U_y = \cap_{j=1}^N U_y^{x_j}$, then $C \subset \mathrm{Int}\,(K^-(y'))\ \forall y' \in U_y$, so that $\mathrm{Int}\,(K^-(\cdot))$ is inner continuous. Similarly one proves the inner continuity of $\mathrm{Int}\,(K^+(\cdot))$. $\Box$

3 Causal Continuity in Morse Spacetimes

Our motivation to consider the relation K^+ is that it provides a framework in which to discuss the causal properties of degenerate spacetimes. As mentioned in the introduction, such degeneracies are a necessary feature of compact, causal Lorentzian topology changing spacetimes. In particular, one can construct a rather general class of such spacetimes which contain only isolated degeneracies. These "Morse spacetimes" were studied in detail in [6, 16, 7] within the Sum-Over-Histories approach to quantum gravity, and can be defined in any topological cobordism, as follows.

Morse spacetimes are defined on a compact cobordism $(\mathcal{M}, V_0, V_1)$, with $\partial\mathcal{M} = V_0 \amalg V_1$, a Riemannian metric h on $\mathcal{M}$ and a Morse function $f : \mathcal{M} \to [0, 1]$, with $f^{-1}(0) = V_0$, $f^{-1}(1) = V_1$ which has r critical points, $\{p_k\}$, in the interior of $\mathcal{M}$. The critical points of a differentiable function f are those where $\partial_\mu f = 0\ \forall \mu$. What characterises a Morse function is that the Hessian $\partial_\mu\partial_\nu f$ of f is an invertible matrix at each of its critical points, which we call Morse points. The Morse index λ_k of the critical point p_k is the number of negative eigenvalues of the matrix $(\partial_\mu\partial_\nu f)(p_k)$. From h, f and an arbitrary

real constant greater than 1, denoted ζ, we can define a *Morse metric* in $\mathcal{M}$:

$$g_{\mu\nu} = (h^{\rho\lambda}\partial_\rho f \partial_\lambda f)h_{\mu\nu} - \zeta \partial_\mu f \partial_\nu f, \tag{7}$$

The vector field $h^{\mu\nu}\partial_\nu f$ is time-like with respect to the metric g, so that we can think of f as a time-function on (M, g). We refer the reader to [7, 6, 16] for a more expository account. The metric (7) is Lorentzian everywhere, except at the critical points $\{p_k\}$ of f. If, as is standard practice in causal analysis, we excise these critical points from the spacetime, the resulting *Morse spacetime*, (M, g), is the globally Lorentzian spacetime induced in $M = \mathcal{M} - \{p_k\}$.

A general cobordism $\mathcal{M}$ can be decomposed into a finite sequence of elementary cobordisms $\mathcal{E}_k$, each containing one critical point. If $\mathcal{E}$ is an elementary cobordism, we will refer to a Morse spacetime (E, g), defined in $E = \mathcal{E} - p$, as an elementary Morse spacetime.

The Morse Lemma tells us that there exists a neighbourhood of a critical point, p, with local coordinates $(x_1, \cdots, x_\lambda, y_1 \cdots, y_{n-\lambda})$, in which the Morse function f takes a canonical quadratic form at any point in this neighbourhood,

$$f(x) = f(p) - \sum_{i=1}^{\lambda} x_i + \sum_{j=1}^{n-\lambda} y_j \tag{8}$$

where λ is the index of the Morse function. In what follows we will refer to these neighbourhood spacetimes as neighbourhood Morse spacetimes, as opposed to the full Morse spacetimes, defined above.

We now proceed to explore the (I^+, K^+) structure for Morse spacetimes both when the degeneracies are excised to give a globally Lorentzian spacetime and when they are left in.

3.1 Morse spacetimes with the critical points excised

In the case of a Morse spacetime with a single critical surface, we can show that when causal continuity fails, then at least one of $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$ is not inner continuous. In other words, the converse of lemma 10.(i) is now true. We know that any causally discontinuous Morse spacetime (M, g) contains a causally discontinuous elementary cobordism (M', g). It suffices to show that then there must be $x, y \in M'$ such that $x \in \mathrm{Int}\,(K^-(y))$ and yet $y \notin \mathrm{Int}\,(K^+(x))$, for this means that $x \in \mathrm{Int}\,(K^-(y))$ and, by the K-convexity of M' (lemma 5), that $y \notin \mathrm{Int}\,(K^+(x))$.

Lemma 12 *For any point x in a Morse spacetime (M, g) where the Morse function f has a single critical value $K^+(x) = \overline{\uparrow I^-(x)}$ and $K^-(x) = \overline{\downarrow I^+(x)}$.*

Proof Let us call $\mathcal{K}^+$ the relation consisting of pairs (x, y) such that $x \in \overline{\downarrow I^+(y)}$. And let $c = f(p)$ be the critical value.

(i) First we check that $x \in \overline{\downarrow I^+(y)}$ if and only if $y \in \overline{\uparrow I^-(x)}$. If the former holds, there are sequences $x_k \to x$ and $y_l \to y$ with $y_l \in I^+(y)\ \forall l$ and $x_k \in I^-(y_l)\ \forall k, l$. Clearly, it suffices to show that $y_l \in \uparrow I^-(x)\ \forall\, l$. For each y_l, there is another y_m and a neighbourhood U_{y_l} of y_l such that $U_{y_l} \subset I^+(y_m)$. Now for any point $x' \in I^-(x)$, there is an x_k with $x' << x_k$. This means that $U_{y_k} \subset I^+(y_m) \subset I^+(x_k) \subset I^+(x') \forall\, x' \in I^-(x)$, as desired. The converse implication is proved analogously.

(ii) The relation $\mathcal{K}^+$ is closed. Let $x_k \in \overline{\downarrow I^+(y_k)}$ with $x_k \to x$ and $y_k \to y$. It suffices to show that any $x' \in I^-(x)$ is in $\downarrow I^+(y)$. Given any such x', we can find neighbourhoods U_x of x and $U_{x'}$ of x' with $U_{x'} \times U_x \subset I^+$. Moreover, there is an integer k_0 such that $x_k \in U_x \quad \forall k > k_0$. Therefore U_x contains points $x'_k \in \downarrow I^+(y_k) \quad \forall k > k_0$. Now for every $y' \in I^+(y)$, there is a y_l with $l > k_0$ to the past of y'. Hence, $U_{x'} \subset I^-(x'_l) \subset I^-(y')$, that is $x' \in \downarrow I^+(y)$.

(iii) The relation $\mathcal{K}^+$ is transitive. It is here that we need the assumption that there is only one critical value. Suppose $x \in \overline{\downarrow I^+(y)}$ and $y \in \overline{\downarrow I^+(z)}$. We wish to show that $x \in \overline{\downarrow I^+(z)}$. We need to consider three cases:

Case 1. $x \in \overline{I^-(y)}$ and $y \in \overline{I^-(z)}$ implies, by claim 1, that $x \in \overline{I^-(z)} \subset \overline{\downarrow I^+(z)}$.

Case 2. $x \in \overline{I^-(y)}$ and $y \in \overline{\downarrow I^+(z)} - \overline{I^-(z)}$. Take a sequence $x_k \to x$ contained in $I^-(y)$ and a sequence $y_k \to y$ in $\downarrow I^+(z)$. For each x_k there is a $y_l >> x_k$ and hence a neighbourhood $U_{x_k} \subset I^-(y_l)$. It follows that for every $z' >> z$, $U_{x_k} \subset I^-(z')$. Thus the sequence $\{x_k\}$ is also contained in $\downarrow I^+(z)$.

Case 3. $x \in \overline{\downarrow I^+(y)} - \overline{I^-(y)}$ and $y \in \overline{I^-(z)}$. We must have $f(x) < c < f(y)$, because the subcritical region is globally hyperbolic. And so is the supercritical region, whence, $y \in J^-(z)$. Consider a sequence $x_k \to x$ with $x_k \in \downarrow I^+(y)\ \forall\, k$, that is, each x_k has a neighbourhood U_{x_k} contained in the past of every $y' >> y$. Take any $z' \in I^+(z)$. It follows from axiom (vi+) in definition 2 that $z' \in I^+(y)$ so that there exists a point $y' \in I^+(y) \cap I^-(z')$. Thus $U_{x_k} \subset I^-(z')\ \forall\, z'\, k$, i.e., $x_k \in \downarrow I^+(z)\ \forall\, k$. □

With this, we can now prove:

Lemma 13 *A Morse spacetime (M, g) is causally continuous if and only if the functions* $\mathrm{Int}\,(K^+(\cdot))$ *and* $\mathrm{Int}\,(K^-(\cdot))$ *are inner continuous.*

Proof We saw in lemma 10 that causal continuity implies inner continuity of $\mathrm{Int}\,(K^+(\cdot))$ and $\mathrm{Int}\,(K^-(\cdot))$. Conversely suppose (M, g) is causally discontinuous and so contains a causally discontinuous elementary subcobordism, M' [8]. This implies the existence of points $x, y \in M'\ x \in \downarrow_{M'} I^+(y, M') - \overline{I^-(y, M')}$. Thus $x \in \mathrm{Int}\,(K^-_{M'}(y))$ by the previous lemma. However, $y \notin \mathrm{Int}\,(K^+_{M'}(x))$. This is because $\mathrm{Int}\,(K^+(x)) \cap M' = \uparrow I^-(x) \cap M'$ and $y \in \overline{\uparrow I^-(x)} - \uparrow I^-(x)$ (otherwise, if y was in $\uparrow I^-(x)$, there would be a sequence

$\{x_k\} \subset I^-(y)$ converging to x and we would have $x \in \overline{I^-(y)}$, a contradiction). lemma 11 $\Box$

3.2 Morse spacetimes with the critical points left in

To incorporate the critical points of a degenerate spacetime $\mathcal{M}$ to the K^+ causal order we choose a natural chronology in $\mathcal{M}$ and find its transitive closure. More exactly, the definition that we propose is:

Definition 4 *Let $(\mathcal{M}, g)$ be Lorentzian everywhere, except for a set $\{p_k\} \subset \mathcal{M}$ of isolated points where g vanishes, $M = \mathcal{M} - \{p_k\}$ be the regular part of the spacetime and $I^+_{\mathcal{M}}$ be the usual chronology defined through timelike curves in $\mathcal{M}$. So $I^+_{\mathcal{M}} = I^+_M$ and the chronological past and future of each degeneracy are empty. We define the relation $K^+_{\mathcal{M}}$ in $\mathcal{M}$ to be the transitive closure of $I^+_{\mathcal{M}}$ in $\mathcal{M} \times \mathcal{M}$.*

The question of uniqueness arises, since one can think of other natural chronologies in a degenerate spacetime. To justify definition 4 we should demonstrate that the transitive closures of the different chronologies are the same.

Strictly, a curve passing through a degeneracy p is neither timelike, null or spacelike, and in this sense we would say that $I^{\pm}(p)$ is empty. On the other hand, there are timelike curves that run off towards p, and it would seem reasonable to say that all the points in such curves belong to $I^-(p)$. This leads us to two natural definitions of chronology in a degenerate spacetime $(\mathcal{M}, g)$:

The first one is $I^+_1 \equiv I^+_M$ is that used in definition 4. This is the chronology defined through timelike curves in $\mathcal{M}$ if we are strict about the conditions on the tangent vector of a causal curve. Because no tangent vector at a degenerate point p_k can be deemed as past or future directed[4] there are no timelike or causal curves through the degeneracies. Thus for each p_k, we have $I^{\pm}_{\mathcal{M}}(p_k) = \emptyset$, i.e., the chronological past and future of each degenerate point is empty.

The second one is I^+_2, which we define by allowing causal curves to go through the degeneracies. To do so, we redefine a future-directed causal curve Γ in $\mathcal{M}$ as a piecewise smooth curve whose tangent vector $\dot{\gamma}^\mu$ obeys $g_{\mu\nu}\dot{\gamma}^\mu\dot{\gamma}^\nu \leq 0$ and is future-directed at all points in $\Gamma \cap M$. Similarly for past-directed causal curves. In this way we define a causal relation J^+_2 for which $J^{\pm}_2(p)$ will in general be non-trivial for a degeneracy p. The chronological relation I^+_2 is then defined as $I^{\pm}_2(q) = \mathrm{Int}\left(J^{\pm}_2(q)\right)$ for every point $q \in \mathcal{M}$.

In figure 3, we illustrate the definitions of I^+_1 and I^+_2 for the trousers spacetime of [6, 7]. One can verify by inspection that K^+_1 and K^+_2 are both equal to J^+_2, as in the

[4] In fact the full geometry $(\mathcal{M}, g)$ does not admit a time orientation, because any vector field in $\mathcal{M}$ will be trivially null at each point p_k. We can still define future and past directed causal curves using the time-orientation available in M.



Figure 3: Dependence of chronology and causality on the definition of timelike and causal curves in the trousers spacetime. The top two figures show $I_1^{\pm}(p)$ and $J_1^{\pm}(q)$, respectively, where q lies on the "light cone" of the degeneracy, p. The bottom two figures show $I_2^{\pm}(p)$ and $J_2^{\pm}(q)$

closed shaded regions in figure 3(b). We show below that in fact, this is true for any Morse spacetime, i.e., that although in general, $I_1^+ \subset I_2^+$, they lead to exactly the same relation K^+ in (M, g).

Lemma 14 *Let $(\mathcal{M}, g)$ be a Morse geometry with the chronologies I_1^+ and I_2^+ in $\mathcal{M}$ defined as above. If K_1^+ and K_2^+ are their respective transitive closures in M, then $K_1^+ = K_2^+$.*

Proof Notice first that $I_1^+ \subseteq I_2^+$ implies $K_1^+ \subseteq K_2^+$. The converse will follow from minimality of K_2^+ if we show that $I_2^+ \subset K_1^+$. What we demonstrate is that in fact $J_2^+(x) \subset K_1^+(x)\ \forall x$. Let f be the Morse function in M. For any $y \in J_2^+(x)$, there is a piecewise smooth path $\gamma : [0,1] \to \mathcal{M}$ with $\dot{\gamma}^\mu$ future-directed null or timelike except possibly for a finite number of degeneracies $x_i = \gamma(t_i)\ i = 1, \ldots r$, through which Γ passes in order of increasing f. Add to the list the points $x_0 = x$ and $x_{r+1} = y$ and let $t_0 = 0$ and $t_{r+1} = 1$. For each j between 0 and r there are sequences $\{t_j^m\} \to t_j$ and $\{s_j^m\} \to t_{j+1}$ with $t_j < t_j^m < s_j^m < t_{j+1}$ such that the associated sequences $y_j^m = \gamma(t_j^m) \to x_j$ and $z_j^m = \gamma(s_j^m) \to x_{j+1}$ consist of regular points in M. Therefore these points must satisfy $(y_j^m, z_j^m) \in J_1^+$, which implies $(x_j, x_{j+1}) \in K_1^+$, since K_1^+ contains J_1^+ and is closed. And hence, by transitivity, $(x, y) \in K_1^+$. $\square$

In order to relate K^+ in spacetimes with and without the degeneracies, one would need a generalised version of Proposition 2. In other words, if $(\mathcal{M}, g)$ is a spacetime with an isolated degeneracy at p and (M, g) the globally Lorentzian spacetime with $M = \mathcal{M} - p$, then we expect that the causal relations $K^+_{\mathcal{M}}$ on $(\mathcal{M}, g)$ and K^+_M on (M, g) should be related as follows,

$$K^+_M = K^+_{\mathcal{M}} \cap M \times M \qquad \text{and} \qquad K^+_{\mathcal{M}} = \overline{K^+_M}, \tag{9}$$

the latter a consequence of the former as in lemma 9.

We do not, however, have a complete proof of this statement, due to a difficulty in proving transitivity when the degenerate point is sandwiched between two regular points of the manifold. Indeed, the proof of Proposition 2 already gives a hint of the nature of the problem, where in proving transitivity, one requires the spacetime to be C^2 at the excised points. Clearly, this is too strong a requirement to be used here.

For the index 0 and n elementary Morse spacetimes, however, since the degenerate point cannot be sandwiched between two regular points of the manifold, there is no problem with transitivity, and one may prove the validity of the condition. From this, and our experience with $2-d$ Morse spacetimes, we suspect that the condition is in fact satisfied for all Morse spacetimes.

We will therefore restrict our attention to those spacetimes satisfying this condition, and investigate its consequences in what follows. We state the condition as:

Condition 1 *For a Morse spacetime (M, g) with causal order K^+_M and $K^+_{\mathcal{M}}$ the extended causal order on $\mathcal{M} = M \cup p$, with p the Morse point, we require $K^+_{\mathcal{M}} \cap M \times M = K^+_M$.*

3.3 Causal continuity in Morse geometries

We would like to know whether the characterisations of causal continuity for globally Lorentzian Morse spacetimes, in proposition 3 and lemma 13 can be extended to Morse spacetimes in which the degeneracies are left in. To define causal continuity in $\mathcal{M}$ we use the extended causal order $K^+_{\mathcal{M}}$. A first possibility is to replace K^+_M by $K^+_{\mathcal{M}}$ in condition (B) of proposition 3 and we propose

Definition 5 *The degenerate spacetime $(\mathcal{M}, g)$, with regular part M, is said to be k-causally continuous if every $x \in M$ satisfies $K^\pm_{\mathcal{M}}(x) = \overline{I^\pm(x)}$.*

Here the closure of $I^+(x)$ is taken in $\mathcal{M}$ instead of M. Under the assumption that Condition 1 holds, we can show that the above definition is simply a restatement of condition (B) in proposition 3, i.e., that it agrees with causal continuity of the globally Lorentzian spacetime (M, g).

Lemma 15 *Let $(\mathcal{M}, g)$ be a spacetime with isolated degeneracies and (M, g) the associated globally Lorentzian spacetime which results from excising the degeneracies. Suppose (M, g) satisfies Condition 1, then $(\mathcal{M}, g)$ is k-causally continuous iff (M, g) is causally continuous.*

Proof

(1) Let (M, g) be causally continuous, i.e., $\forall x \in M$, $K^+_M(x) = \overline{I^+(x)}^M$, where the superscript ^{M} indicates that the closure is taken in M. We want to prove that a point $y \in K_{\mathcal{M}}(x)$ must also satisfy $y \in \overline{I^+(x)}^{\mathcal{M}}$.

a) If $y \in M$, then Condition 1 says that $y \in K^+_M(x) = \overline{I^+(x)}^M \subset \overline{I^+(x)}^{\mathcal{M}}$.

b) If $y \notin M$, that is if y is one of the degeneracies. Then from lemma 3 we can find a sequence of regular points $y_k \to y$ such that $y_k \in K^+_{\mathcal{M}}(x)$. It follows from Condition 1 that $y_k \in K^+_M(x) = \overline{I^+(x)}^M \subset \overline{I^+(x)}^{\mathcal{M}}$. And therefore $y \in \overline{I^+(x)}^{\mathcal{M}}$, as required.

(2) Let $(\mathcal{M}, g)$ be k-causally continuous. Suppose $y \in K^+_M(x)$, then $y \in K^+_{\mathcal{M}}(x) = \overline{I^+(x)}^{\mathcal{M}}$. Now, since $y \in M$, a sequence $y_k \to y$ must exist with $y_k \in I^+(x)$ and such that $y_k \in M$ $\forall k$. Therefore $y \in \overline{I^+(x)}^M$, which means (M, g) is causally continuous. □

Note that part (2) of the proof does not depend on Condition 1 and therefore holds in general. In other words, if (M, g) is causally discontinuous then $(\mathcal{M}, g)$ is k-causally discontinuous. From [8] we know that (M, g) is causally discontinuous if it contains index 1 or $n - 1$ Morse points, which means that it is k-causally discontinuous as well.

To extend definition 5 to the degeneracies of a Morse spacetime, we first have to choose an appropriate definition of chronology $I^{\pm}(p)$ for each Morse point p. By contrast, conditions (D) and (E) in the previous section are expressed in terms of K^+ alone, so that one may expect them to extend directly to the whole degenerate spacetime by simply replacing K^+_M by $K^+_{\mathcal{M}}$. Are these conditions equivalent to causal continuity of the punctured Morse spacetimes?

The extension of condition (D) is: $\mathrm{Int}\,(K^+_{\mathcal{M}}(\cdot))$ and $\mathrm{Int}\,(K^-_{\mathcal{M}}(\cdot))$ are inner continuous as functions on $\mathcal{M}$. If the spacetime were to satisfy the condition, $K^+_{\mathcal{M}} \cap M \times M = K^+_M$ (Condition 1), then inner discontinuity of $\mathrm{Int}\,(K^+_M(\cdot))$ would imply that of $\mathrm{Int}\,(K^+_{\mathcal{M}}(\cdot))$ (similarly for $\mathrm{Int}\,(K^-(\cdot))$.) Thus causal discontinuity of the Morse spacetime implies inner discontinuity of the extended $\mathrm{Int}\,(K^+_{\mathcal{M}}(\cdot))$ or $\mathrm{Int}\,(K^-_{\mathcal{M}}(\cdot))$. However, the converse is not true: causal continuity does not always imply inner continuity of the extended $\mathrm{Int}\,(K^+)$ and $\mathrm{Int}\,(K^-)$.

A counterexample that illustrates this is a two-dimensional yarmulke, *i.e.* a Morse geometry with $n = 2$, $\lambda = 0$, corresponding to the topology change $\emptyset \to S^1$. Let x, y be the coordinates in which the Morse function takes the form $f = x^2 + y^2$, let $\mathcal{M} = \{(x, y) : x^2 + y^2 < 1\}$, let the Riemannian metric h be given by the interval $ds^2_R = dx^2 - 2dxdy + 2dy^2$ in these coordinates and set $\zeta = 2$. Studying the null-

equations of the Morse metric g constructed from f, h and ζ, we obtain the light-cone structure drawn in figure 4. From [8] we know that the Morse spacetime $(\mathcal{M}-p, g)$ is causally continuous. However, it is easy to see that $\mathrm{Int}\,(K^+_{\mathcal{M}}(\cdot))$ is not inner continuous. Let A be the closed region shown in figure 4 bounded by the lines $y=0$ and $y=m_3x$. Now, although $K^+_{\mathcal{M}}(p)=\mathcal{M}$, for any $x\in A$, $K^+_{\mathcal{M}}(x)\subset A$ [5].

Thus, although the spacetime $(\mathcal{M}, g)$ is causally continuous, $\mathrm{Int}\,(K^+_{\mathcal{M}}(\cdot))$ is not inner continuous at p. It is therefore perhaps inappropriate to use the extended version of condition (D) as the definition of k-causal continuity when the degenerate points are included.



Figure 4: The yarmulke neighbourhood geometry constructed from the Riemannian metric with interval $ds^2_R = dx^2 - 2dxdy + 2dy^2$. Shown are the four null straight lines through the origin, with gradients m_1, $m_2=0$, m_3 and $m_4=\infty$. The straight lines $y=mx$ are spacelike for gradients m between m_1 and m_2 and between m_3 and m_4; and they are timelike for the remaining values of m. Representative lightcones are shown for points on the null radii, with future direction being outwards from the morse point p. No point in this spacetime contains a whole neighbourhood of p in its past and the causal future of any point in the region A, say, is contained in A.

For condition (E), the extension in terms of $K^+_{\mathcal{M}}$ on $\mathcal{M}$ is: for any $x\in\mathcal{M}$ and $K_{\mathcal{M}}$-convex neighbourhood N of x, $\mathrm{Int}\,(K^+_N(x))$ and $\mathrm{Int}\,(K^-_N(x))$ are connected. We know of no counterexample to the possibility that condition (E) for $\mathcal{M}$ is equivalent to causal continuity of the spacetime with the degeneracies excised, but neither do we have a proof.

[5] Note however, that if, in the spirit of definition 5, we had only defined condition (D) to hold at the regular points of $\mathcal{M}$, then the above yarmulke would in fact satisfy it. However, our interest is in a definition which doesn't single out degenerate points

4 Discussion

We have presented preliminary results on the causal structure, and causal continuity in particular, in terms of the natural and robust causal relation K^+ on spacetimes with isolated degeneracies. The case of most interest to us is that of topology changing Morse spacetimes. To make further progress, it is necessary for us to prove that causal continuity of a degenerate spacetime can be characterised entirely in terms of K^+.

Ultimately, topology changing processes can only be analysed with rigour in a theory of quantum gravity. Any seemingly ad hoc rules governing topology change should be justified by appealing to the more basic theory. Underlying our analysis, then, is the idea that perhaps the causal order of a spacetime is its most fundamental aspect. This is the basis of the causal set hypothesis [12] that the fundamental discrete (quantum) substructure of spacetime is described by a causal set.

The question thus arises: can any sense be made at the discrete level of a continuum notion like causal continuity? In a sense we have been working towards this in this paper — framing conditions in terms of the K^+ relation alone and independently of I^+, for example, could be seen as moving in this direction, since a causal set possesses a causal relation but not a chronological one.

Basic to the causal set hypothesis is the notion of "manifold-like" causal sets which can be obtained by sprinkling points at random, with a fixed density, into a Lorentzian manifold and using the spacetime causal order to define the partial order on the set of sprinkled points. We can pursue the question of whether causal continuity makes sense in a causal set by asking what a causal set that arises from sprinkling into causally discontinuous spacetimes looks like. A preliminary investigation suggests that causal discontinuity might manifest itself in the causal set in the existence of a point x in the set and a single link from x, the severing of which would result in a big change in the causal future or past of x in the causal set. What is meant by big change and whether this can be made more concrete is left for future work.

5 Acknowledgments

We would like to thank Rafael Sorkin and Eric Woolgar for valuable discussions.

References

[1] R.D.Sorkin and E.Woolgar. A causal order for spacetimes with c^0 lorentzian metrics: Proof of compactness of the space of causal curves. *Class.Quant.Grav.*, 13:1971–1994, 1996.

[2] S.W. Hawking and G.F.R. Ellis. *The large scale structure of space-time*. Cambridge University Press, Cambridge, 1973.

[3] R.Penrose. *Techniques of differential topology in relativity*. Society for Industrial and Applied Mathematics, Pennsylvania, 1973.

[4] R.P.Geroch. Topology in general relativity. *J. Math. Phys*, 8:782, 1967.

[5] R.D. Sorkin. On topology change and monopole creation. *Phys. Rev.*, D33:978, 1986.

[6] Dowker and Surya. Topology change and causal continuity. *Phys. Rev.*, D58:124019, 1998.

[7] Borde, Dowker, Garcia, Sorkin, and Surya. Causal continuity in degenerate space-times. *Class. Quantum Grav.*, 16:3457–3481, 1999.

[8] Dowker, Garcia, and Surya. Morse index and causal continuity: A criterion for topology change in quantum gravity, 1999. Accepted for publication in Class. Quant. Grav.

[9] R.P.Geroch, E.H.Kronheimer, and R.Penrose. *Proc.Roy.Soc.Lond.A*, 327:545, 1972.

[10] Hawking and Sachs. Causally continuous spacetimes. *Comm. Math. Phys*, 35:287–296, 1974.

[11] E.H.Kronheimer and R.Penrose. *Proc.Cam.Phyl.Soc*, 63:481, 1967.

[12] Bombelli, Lee, Meyer, and Sorkin. Space-time as a causal set. *Phys.Rev.Lett*, 59:521, 1987.

[13] H.J. Seifert. *Gen. Rel. Grav.*, 1:247, 1971.

[14] J.Kelley. *General Topology*. van Nostrand, Toronto, 1955.

[15] R.D. Sorkin, 1999. Private Communication.

[16] H.F. Dowker and R.S. Garcia. A handlebody calculus for topology change. *Class. Quantum Grav.*, 15:1859, 1998. gr-qc/9711042.